

Time Evolution of Thermal Photon Elliptic Flow

Rupa Chatterjee, Dinesh K. Srivastava

Variable Energy Cyclotron Centre, 1/AF, Bidhan Nagar, Kolkata 700 064, India

Ulrich Heinz

Physics Department, The Ohio State University, Columbus, OH 43210, USA

Abstract

Elliptic flow of thermal photons has a great potential to explore the early time dynamics and evolution of Quark Gluon Plasma. p_T dependent temporal contours for photon spectra and elliptic flow from quark matter and hadronic phases, as well as the p_T integrated results show gradual build-up of flow with time and relative contributions from different phases to that very clearly. Unlike hadrons, photon flow is quite sensitive to the initial thermalization time τ_0 , and its value can be estimated with the experimental determination of v_2 .

Key words: Photons, elliptic flow, time evolution, τ_0 etc.

PACS:

1. Introduction

The observation of large anisotropic flow or in particular elliptic flow of different hadronic species at the Relativistic Heavy Ion Collider (RHIC) at Brookhaven National Lab, New York, provides a strong confirmation of the formation of QGP. Photons are known to probe the properties of QGP in a relatively clean manner compared to hadrons. Unlike hadrons, photons are emitted from every stage of the expanding system and also carry undistorted information from the production point to the detector. Elliptic flow is known to provide information about early thermalization and collectivity in the hot and dense matter. The flow parameter v_2 is quantified as the 2nd Fourier co-efficient of the particle distribution in the transverse momentum plane as (at mid-rapidity and for collisions of same type of nuclei only even cosine terms survive in the series below):

$$\frac{dN(b)}{p_T dp_T dy d\phi} = \frac{dN(b)}{2\pi p_T dp_T dy} [1 + 2v_2(p_T, b) \cos(2\phi) + 2v_4(p_T, b) \cos(4\phi) + \dots] \quad (1)$$




Fig. 1. Left panel: v_2 for thermal photons along with contributions from QM and HM. Right panel: p_T integrated v_2 results from different phases as a function of τ .

We have shown that elliptic flow of thermal photons for 200A GeV Au+Au collisions at RHIC (using ideal hydrodynamic model) reflects the anisotropies of the early partonic phase at large values of transverse momentum p_T [1]. In a recent calculation of photon v_2 (considering prompt and jet photons along with thermal), it is shown that the thermal v_2 dominates over others for $p_T < 5$ GeV [2].

2. Time evolution of photon v_2

The time evolution of photon spectra and v_2 is studied for impact parameter $b = 7$ fm and $\tau_0 = 0.2$ fm considering the same initial conditions as given in Ref. [1]. Rates of photon production from quark matter and hot hadronic gas are taken from Arnold *et al.* [3] and Turbide *et al.* [4] respectively, which provide good description of the single photon data at RHIC as well as at SPS energies. The p_T dependent elliptic flow parameter for thermal photons along with contributions to v_2 from quark matter (QM) and hadronic matter (HM) phases are shown separately in left panel of Fig.1. In the right panel of same figure, the p_T integrated v_2 from different phases as a function of proper time τ and normalized with $v_2(\tau_f)$ from respective phases are shown, where τ_f (≈ 12 fm) is the time when freeze-out is completed [5]. As shown in the figure, the $v_2(QM)$ saturates early within a time period of 5 fm. The $v_2(HM)$ is small at the beginning and saturates much later compared to $v_2(QM)$. Contours of p_T dependent spectra and v_2 at various values of τ from QM, HM and the sum of the two phases are shown explicitly in Fig. 2. At large p_T (> 3 GeV), most of the photons from QM are emitted very early within a time period of 2 fm. However, the $v_2(QM)$ is not that strong at early times and it gradually builds-up as shown in the left panel of Fig. 2. Photons from HM and their v_2 , both are very small at the beginning and become significant only after 4-5 fm. The sum spectra and v_2 show clear QM dominance at early times or large p_T and the HM contribution is substantial only for lower p_T or later times.

3. τ_0 from elliptic flow of thermal photons

The final $v_2(p_T)$ for photons is quite sensitive to the initial thermalization time τ_0 for a fixed value of particle multiplicity, and the value of τ_0 can thus be estimated with the help of experimental results [6]. The differential elliptic flow $v_2(p_T)$ for different τ_0 ranging from 0.2 to 1.0 fm at $b = 6$ fm, are shown in left panel of Fig. 3. For smaller τ_0 ,




Fig. 2. p_T spectra [upper panel] and elliptic flow [lower panel] of thermal photons for different values of τ from QM, HM and the sum of the two phases.




Fig. 3. $v_2(p_T)$ for thermal photons [left panel] and π mesons [right panel] at different τ_0 .

the number of photons from QM increases compared to larger τ_0 at high p_T . As $v_2(QM)$ is smaller at earlier times and $v_2(HM)$ does not change significantly with changing τ_0 , the total v_2 decreases with smaller τ_0 . As hadrons are emitted from freeze-out surface, their v_2 is affected only marginally with changing values of τ_0 (right panel of Fig. 3).

In conclusion, the time evolution results for photon spectra and elliptic flow show gradual build-up of the flow parameter along with relative contributions from different phases with time and at different p_T . Photon flow is quite sensitive to the value of τ_0 , which can be estimated by experimental determination of the flow parameter.

References

- [1] R. Chatterjee, E. Frodermann, U. Heinz, and D. K. Srivastava, Phys. Rev. Lett. **96**, 202302 (2006).
- [2] S. Turbide, C. Gale, E. Frodermann, U. Heinz. Phys. Rev. C **77**, 024909 (2008).
- [3] P. Arnold, G. D. Moore, and L. G. Yaffe, JHEP **0112**, 009 (2001).
- [4] S. Turbide, R. Rapp, and C. Gale, Phys. Rev. C **69**, 014903 (2004).
- [5] R. Chatterjee, D. K. Srivastava, and U. Heinz. [in preparation].

- [6] R. Chatterjee and D. K. Srivastava, e-Print: arXiv:0809.0548 [nucl-th].